



 **Safety**
2010
World
Conference

The FARE

A new way to express Falls Risk among older persons including physical activity as a measure of Exposure

GJ Wijlhuizen, AMJ Chorus, M Hopman-Rock
TNO, Leiden, The Netherlands

TNO




Falls among older persons (55+) in the Netherlands

- Each year:
 - 88.000 Treatments at ER
 - 32.000 Hospital admissions (78% fractures)
 - 1.800 Fatalities
 - High falls injury risk among older persons 75+
 - About 30% older persons fall at least once
 - Direct medical costs: 550 million Euros (700 million USD)

General assumption

- Persons will not fall if:
 - their capability to control balance
 - is greater than
 - the demands put on it.

The occurrence of a fall; capability-demand







Safety of system or person


- Product of:
 - Probability of having an accident/injury given a unit of exposure
 - Observed level of exposure

(Hauer, Accid Anal Prev, 1982; Qin et al, Accid Anal Prev, 2004)


Concept of exposure related to falls

- Exposure is zero if a person does not move (PA level =0)
- Exposure increases if PA level $\gg 0$

Exposure as a precondition for falls



Falls research models do not take exposure into account




Is exposure important in falls research?

- No, if:
 - There are no differences in exposure among older persons
 - If exposure measures show no relation with falls


Any differences in exposure among older persons?

- Subjects were asked for both summer and winter about:
- The number of days during an average week at which they were physically active for at least 30 minutes at a moderate level (at least as heavy as brisk walking or bicycling).



Number of physical active days per week by age (N= 21.020) IPAN data 2000-2005




Exposure by balance difficulty among community dwelling older persons (70+), N=704



Distributions of falls at home (n=305) (top) and physically active person-hours at home (n=459) among community dwelling older persons 65+



Distribution of ratio falls/ per active person-hours at home (Spearman cor = .89)



Wijlhuizen et al, Prev Med, 2008

Conclusions

- There are large differences in exposure (PA) among older persons
- Exposure (PA) measure shows strong relation with falls
- Therefore:
- We should take exposure into account in falls research

Wijlhuizen et al, Int J Inj Control Saf Promot, 2007


Wijlhuizen et al Prev Med, 2007, 2008a,b, 2010.

The Falls risk by Exposure (FARE)


- Common expression of falls risk:
 - Number of falls/ 1000 person-years

- The FARE:
 - Number of falls/ 1000 physically active person-days

Falls injury risk ratio by age expressed by Incidence/FARE (N= 21.020) IPAN data 2000-2005




Falls risk ratio by level of Balance control difficulty expressed by Incidence/FARE (N=704)



Wijlhuizen et al, Prev Med, 2010

Final conclusions

- We should take exposure into account in falls research
 - Because:
 - There are large differences in exposure (PA) among older persons
 - Exposure (PA) measure shows strong relation with falls
 - Actual falls risk is generally strongly underestimated compared to the FARE
- We do not have an exposure measure which is agreed upon



 **Safety
2010
World
Conference**