

MIXED TREATMENT COMPARISONS TO EVALUATE THE EFFECTIVENESS OF STRATEGIES FOR PREVENTING FIRE RELATED INJURIES IN CHILDREN WITHIN THE HOME

Pedro Saramago Goncalves

PhD Student, Centre for Health Economics, University of York


In collaboration with Professor Alex Sutton, Dr Nicola Cooper, Paula Dhiman: University of Leicester & Professor Denise Kendrick: University of Nottingham. On behalf of the 'Keeping Children Safe' project.

OBJECTIVES

- To identify the most effective (i.e. “best”) strategy for increasing the ownership of fire-related safety equipment in households (smoke alarms, fire extinguishers, fire guards, safe storage of matches/lighters, fire escape plans).

METHODS: Mixed Treatment Comparisons

- MTC methods (a generalisation of meta-analysis methods) allow comparisons of strategies not directly assessed within any individual primary study.



- MTC uses the totality of the evidence to define a network of evidence

STAGES IN MTC

1

- Find the evidence

2

- Define the network


3

- Synthesise the evidence

1. FINDING THE EVIDENCE

- Updated Cochrane systematic review of home safety education +/- safety equipment for the prevention of home injuries in childhood and meta-analyses of studies comparing intervention to usual care (*Kendrick et al 2007*)
- Systematic review of reviews with data extraction from relevant primary studies not in meta-analyses in Cochrane review (e.g. smoke alarm education vs. smoke alarm education + free fitted alarm)

2. DEFINING THE NETWORK



E.g.
Interventions
for increasing
ownership of
functioning
smoke alarms

3. SYNTHESISING THE EVIDENCE

- **Mixed treatment random effects model** with a binary outcome (*have functioning equipment / do not have functioning equipment*)
- **Adjusted for clustering** in cluster trials that did not in original analysis (ICCs estimated from individual studies/literature)
- **Outcomes:**
 - Odds ratio of each intervention compared to one another
 - Probability of each intervention being the best

RESULTS: *Probability intervention is the “best”*

Probability	Smoke alarms	Fire extinguishers	Fire guards	Matches/lighters safe storage	Fire escape plans
Usual care	0.000	0.017	0.008	0.017	0.030
Education	0.000		0.126	0.025	0.144
Education + equipment	0.210		0.630		
Education + equipment (not related)		0.063			0.477
Education + equipment + home inspection	0.160	0.752	0.126	0.038	
Education + equipment (not related) + home inspection		0.107			
Education + equipment + home inspection + fitting			0.110		
Education + equipment + fitting	0.120			0.920	
Education + home inspection	0.510	0.030			
Community campaigns		0.031			0.349

RESULTS: *Odds Ratios versus standard care*

<i>OR versus standard care</i>	Smoke alarms	Fire extinguishers	Fire guards	Matches/lighters (safe storage)	Fire escape plans
Education	1.049 (0.14 to 3.944)		3.136 (0.081 to 15.62)	3.804 (0.09 to 14.2)	4.66 (0.163 to 28.34)
Education + equipment	4.901 (0.821 to 16.57)		10.84 (0.326 to 64.6)		
Education + equipment (not related)		1.773 (0.081 to 8.977)			17.82 (0.093 to 94.07)
Education + equipment + home inspection	5.862 (1.182 to 20.5)	11.06 (0.465 to 66.94)	2.403 (0.092 to 14.25)	1.244 (0.225 to 4.175)	
Education + equipment (not related) + home inspection		2.525 (0.036 to 19.09)			
Education + equipment + fitting	3.449 (0.537 to 12.24)		2.795 (0.061 to 13.72)	4127 (0.579 to 9299)	
Education + home inspection	14.21 (0.82 to 73.89)	1.215 (0.07 to 5.706)			
Community campaigns		1.097 (0.024 to 5.262)			5.268 (0.167 to 25.97)

DISCUSSION

- Mixed Treatment Comparison Methods allow a coherent analysis of the totality of the evidence
- Major assumption of the approach:
 - The intervention effect estimated by the *BC* trials would be the same as the intervention effect estimated by the *AC* and *AB* trials if they had included *B* and *C* arms etc.
 - Violated if factors interact with intervention effects and these factors imbalanced across interventions (e.g. social group of study participants etc.)
- Quality/validity of studies ignored
 - As for standard meta-analysis, difficult to incorporate
- Small number of studies for most comparisons
- Results from these analyses are to be used in decision modelling to ascertain cost-effectiveness of the different interventions

CONCLUSIONS

- Results from the MTC show that more intensive interventions are generally estimated to be more effective (i.e. with the best strategies usually offering free or low cost equipment)
- The best strategies varies across equipment types
 - may partly be due to limitations in the data
- Further studies needed to increase precision of estimates and distinguish most effective components of interventions

References and contact details

- **References:**

Kendrick, D., Coupland, C., Mulvaney, C., Simpson, J. Smith, S.J., Sutton, A., Watson, M., Woods, A. Home safety education and provision of safety equipment for injury prevention. Cochrane Database of Systematic Reviews 2007, CD005014.

Caldwell DM, Gibb DM, Ades AE. Validity of indirect comparisons in meta-analysis. The Lancet. 2007;369:270-1.

Sutton, A.J., Ades, A.E., Cooper, N.J., Abrams, K.R. Use of indirect and mixed treatment comparisons for technology assessment. Pharmacoeconomics 2008: 26:753-767.

- **Contact details:**

prsg500@york.ac.uk

This study was funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research funding scheme (RP-PG-0407-10231). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.